REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments reparding this burden estimate or any other ascect of this collection of information, including suggest one for requiring this ourgen to Wishington headquarters for recommendation Operations and Reports, 1215 Lefferson Davis Highway, Suite 1204, Arlington, VA 22202-4322, and to the Office of Management and Budget, Piperwork Reduction Project (0704-0188), Washington, OC 20503.

1. AGENCY USE DNLY (Leave blank)

2. REPORT DATE 9/23/96 3. REPORT TYPE AND DATES COVERED Final

4. TITLE AND SUBTITLE

Ligands for stretch activated ion channels

5. FUNDING NUMBERS
DAAL032-92-G-0014

6. AUTHOR(S)

Frederick Sachs

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

State University of New York at Buffalo Sponsored Programs Administration UB Commons, Suite 211 Amnerst, NY 14228 8. PERFORMING ORGANIZATION REPORT NUMBER

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

U.S.Army Research Office P.O.Box 12211

Research Triangle Park, NC 27709-2211

10. SPONSORING / MONITORING AGENCY REPORT NUMBER

ARO 28775.7-LS

11. SUPPLEMENTARY NOTE:

19961021 159

12a. DISTRIBUTION AVAILABLE STATEMENT

Approved for public release; distribution unlimited.

13. ABSTRACT

During tenure of the grant we have made tremendous strides in finding active ligands for the mechanosensitive channels. We developed a simple screen for the toxins based upon hypotonic swelling of GH3 clonal neurons which produced increases in internal Ca^{2+} levels that could be measured using Fura-2. Addition of active venoms would lead to a decrease in Ca^{2+} levels following swelling. Screening a variety of spider and scorpion venoms, we found that none of the scorpions tested (\approx 12) but one of the \approx 8 spiders tested was able to block volume activated Ca^{2+} uptake. The raw venom also blocked stretch activated ion channels in *Xenopus* oocytes, chick heart cells and GH3 cells, and whole cell mechanical currents in chick heart cells.

14. SUBJECT TERMS

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION 19. SECURITY CLASSIFICATION 20. LIMITATION OF ABSTRACT OF REPORT OF THIS PAGE OF ABSTRACT Unclassified Unclassified Unclassified ULL

USARO Final Report DAAL032-92-G-0014, "Ligands for stretch activated ion channels"

Abstract:

During tenure of the grant we have made tremendous strides in finding active ligands for the mechanosensitive channels. We developed a simple screen for the toxins based upon hypotonic swelling of GH3 clonal neurons which produced increases in internal Ca^{2+} levels that could be measured using Fura-2. Addition of active venoms would lead to a decrease in Ca^{2+} levels following swelling. Screening a variety of spider and scorpion venoms, we found that none of the scorpions tested (\approx 12) but one of the \approx 8 spiders tested was able to block volume activated Ca^{2+} uptake. The raw venom also blocked stretch activated ion channels in *Xenopus* oocytes, chick heart cells and GH3 cells, and whole cell mechanical currents in chick heart cells.

Publications arising from USARO support on this grant:

Chen, Y., S.M. Simasko, J. Niggel, W.J. Sigurdson, and F. Sachs (1996) Ca²⁺ uptake in GH3 cells during hypotonic swelling: the sensory role of stretch-activated ion channels. Am.J.Physiol. 270:C1790-C1798.

Diamond, S.L., F. Sachs, and W.J. Sigurdson (1994) The mechanically induced calcium mobilization in cultured endothelial cells is dependent on actin and phopholipase.

Arteriosclerosis and Thrombosis 14:2000-2009.

Hu, H. and F. Sachs (1993) Effects of mechanical stimulation on embryonic chick heart cells. Upstate NY Cardiac Electrophysiology Society (Abstract)

Hu, H. and F. Sachs (1994a) Effects of mechanical stimulation on embryonic chick heart cells. Biophys.J. 66:A170(Abstract)

Hu, H. and F. Sachs (1994b) Characterizing whole-cell mechanosensitive currents in chick heart. The Physiologist 37:A9(Abstract)

Hu, H. and F. Sachs (1994c) Whole cell mechanosensitive current and its correlation with mechanosensitive channels in chick heart. Upstate NY Cardiac Electrophysiology Society (Abstract)

Hu, H. and F. Sachs (1995) Whole cell mechanosensitive currents in acutely isolated chick heart cells: correlation with mechanosensitive channels. Biophys.J. 68:A393(Abstract)

Hu, H. and F. Sachs (1996a) Mechanically activated currents in chick heart cells. J.Membr.Biol. 154: (in press)

Hu, H. and F. Sachs (1996b) Single-channel and whole-cell studies of mechanosensitive currents in chick heart. Biophysical J. 70:A347(Abstract)

Izu, Y.C., S. Simasko, and F. Sachs (1993) Calcium pathways involved in the hypotonicity-induced calcium increase in GH3 cells. FASEB Journal 7(4):A891(Abstract)

Izu, Y.C. and F. Sachs (1994) Hypotonic cell swelling induced intracellular calcium increase and whole cell currents in GH3 cells: a response to mechanical stimulus and a mechanism for volume regulation. Upstate NY Cardiac Electrophysiology Society (Abstract)

Nazir, S.A., D.J. Dick, F. Sachs, and M.J. Lab (1995) Effects of *G. spatulata* venom, a novel stretch-activated channel blocker, in a model of stretch-induced ventricular fibrillation in the isolated heart. Circ. 292:I-641, #3076(Abstract)

Niggel, J., H. Hu, W.J. Sigurdson, C. Bowman, and F. Sachs (1996) *Grammostola spatulata* venom blocks mechanical transduction in GH3 neurons, *Xenopus* oocytes and chick heart cells. Biophysical J. 70:A347(Abstract)

Ruknudin, A., F. Sachs, and J.O. Bustamante (1993) Stretch-activated ion channels in tissue-cultured chick heart. Am.J.Physiol. 264:H960-H972.

Ruknudin, A.M., H. Hu, and F. Sachs (1994) Extracellular ATP modulates stretch-activated channels in chick heart cells. The Physiologist 37:A6(Abstract)

Sachs, F. (1993a) Ligands for stretch activated ion channels. 6th Int.Symp.Nondestructive Eval.Mat. 124(Abstract)

Sachs, F. (1993b) Stretch activated whole cell currents - whole cell correlates. Trans.Bioelec.Repair Growth Soc.(Abstract)

Sachs, F. (1994) Modeling mechanical-electrical transduction in the heart. In V.C. Mow, F. Guliak, R. Tran-Son-Tray, and R.M. Hochmuth (eds): Cell Mechanics and Cellular Engineering. New York: Springer Verlag, pp. 308-328.

Sachs, F. (1995a) A low drift micropipette holder. Eur.J.Physiol. 429:434-435.(Abstract)

Sachs, F. (1995b) Mechanically sensitive ion channels: biological models for nanoscale stress sensors. Nondestr. Charac. Mater. 6 621-628.

Sachs, F., F. Qin, and P. Palade (1995) Models of Ca²⁺ release adaptation. Science 267:2010-2011.

Sachs, F. and H. Hu (1993) Stretch sensitive whole cell currents are alive and well. Ann.Biomed.Eng. 21:39(Abstract)

Sigurdson, W.J., F. Sachs, and S.L. Diamond (1993) Mechanical perturbation of cultured human endothelial cells causes rapid increases of intracellular calcium. Am.J.Physiol. *264*:H1745-H1752.

Yang, X.C., and F. Sachs (1993) Mechanically sensitive, non-selective, cation channels. In D. Siemen and J. Hescheler (eds): Non-selective ion channels. Heidelberg: Springer-Verlag, pp. 79-92.

Zabel, M., B.S. Koller, F. Sachs, and M.R. Franz (1996) Stretch-induced changes in the isolated heart: importance of the timing of stretch and implications for stretch-activated ion channels. Cardiovas.Res. 32:120-130.

Izu, Y.C. (1994) A study of the sensory mechanism of cell volume regulation. PhD. Thesis, Biophysical Sciences, SUNY, Buffalo, NY.

Inventions:

Sachs, F., Bowman, C.B., Y. Chen-Izu. High Affinity Ligand for Mechanosensitive Ion Channels. SUNY disclosure submitted. Patent application submitted.

Degrees granted with grant support:

Y. Chen-Izu, 1994, SUNY Biophysics.

H. Hu, 1996, SUNY, Biophysics